A Guide to KEYTRUDA® (pembrolizumab) and How to Manage Immune-Mediated Adverse Events

These slides are provided to UK healthcare professionals as a data resource for professional education.

To ensure compliance with all relevant codes and regulations, these slides are provided in PDF format and must not be amended. This content is intended to be viewed online and it is not intended to be printed.

Prescribing information can be found via the 'UK PI' button on each slide. This link will direct you to a third party website.

Please consult the SmPC and risk minimisation materials for further information to minimise the risks associated with the use of the medicine before making any prescribing decisions.



Merck Sharp & Dohme (UK) Limited Registered Office: 120 Moorgate, London EC2M 6UR, United Kingdom Registered in England No. 233687





Adverse events should be reported. Reporting forms and information can be found at <u>https://yellowcard.mhra.gov.uk/</u> or search for MHRA Yellow Card in the Google Play or Apple App Store. Adverse events should also be reported to MSD (Tel: 0208 1548000). By clicking this link, you will be redirected to the MHRA website.



How to use this immune-mediated adverse events guide

- This is an interactive PDF developed as a guide to KEYTRUDA[®] (pembrolizumab) and how to manage immune-mediated adverse events (imAEs)
- Please use the below icons to navigate around this guide:



Links, abbreviations and references

Links to external sites

- The URL links in this slide deck will redirect you to third-party websites. Please note that:
 - MSD does not review or control the content of any third-party website
 - MSD does not endorse and is not responsible for the accuracy, content, practices or standards of any third-party sources

References and abbreviations

• References, and definitions of all abbreviations used in this deck can be found at the end of the presentation





KEYTRUDA[®] (pembrolizumab) indications¹

<u>Melanoma</u>

- **KEYTRUDA** as monotherapy is indicated for the treatment of adults and adolescents aged 12 years and older with advanced (unresectable or metastatic) melanoma.
- KEYTRUDA as monotherapy is indicated for the adjuvant treatment of adults and adolescents aged 12 years and older with Stage IIB, IIC or III melanoma and who have undergone complete resection.

Non-small cell lung carcinoma (NSCLC)

- **KEYTRUDA**, in combination with platinum-containing chemotherapy as neoadjuvant treatment, and then continued as monotherapy as adjuvant treatment, is indicated for the treatment of resectable non-small cell lung carcinoma at high risk of recurrence in adults
- **KEYTRUDA** as monotherapy is indicated for the adjuvant treatment of adults with non-small cell lung carcinoma who are at high risk of recurrence following complete resection and platinum-based chemotherapy.
- **KEYTRUDA** as monotherapy is indicated for the first-line treatment of metastatic non-small cell lung carcinoma in adults whose tumours express PD-L1 with a ≥ 50% tumour proportion score (TPS) with no EGFR or ALK positive tumour mutations.
- **KEYTRUDA**, in combination with pemetrexed and platinum chemotherapy, is indicated for the first-line treatment of metastatic non-squamous non-small cell lung carcinoma in adults whose tumours have no EGFR or ALK positive mutations.
- **KEYTRUDA**, in combination with carboplatin and either paclitaxel or nab-paclitaxel, is indicated for the first-line treatment of metastatic squamous non-small cell lung carcinoma in adults.
- KEYTRUDA as monotherapy is indicated for the treatment of locally advanced or metastatic non-small cell lung carcinoma in adults whose tumours express PD-L1 with a ≥ 1% TPS and who have received at least one prior chemotherapy regimen. Patients with EGFR or ALK positive tumour mutations should also have received targeted therapy before receiving KEYTRUDA.

Classical Hodgkin lymphoma (cHL)

• **KEYTRUDA** as monotherapy is indicated for the treatment of adult and paediatric patients aged 3 years and older with relapsed or refractory classical Hodgkin lymphoma who have failed autologous stem cell transplant (ASCT) or following at least two prior therapies when ASCT is not a treatment option.





KEYTRUDA[®] (pembrolizumab) indications¹

Urothelial carcinoma

- KEYTRUDA, in combination with enfortumab vedotin, is indicated for the first-line treatment of unresectable or metastatic urothelial carcinoma in adults.
- **KEYTRUDA** as monotherapy is indicated for the treatment of locally advanced or metastatic urothelial carcinoma in adults who have received prior platinum-containing chemotherapy.
- **KEYTRUDA** as monotherapy is indicated for the treatment of locally advanced or metastatic urothelial carcinoma in adults who are not eligible for cisplatin-containing chemotherapy and whose tumours express PD-L1 with a combined positive score (CPS) ≥ 10.

Head and neck squamous cell carcinoma (HNSCC)

- **KEYTRUDA**, as monotherapy or in combination with platinum and 5-fluorouracil (5-FU) chemotherapy, is indicated for the first-line treatment of metastatic or unresectable recurrent head and neck squamous cell carcinoma in adults whose tumours express PD-L1 with a CPS ≥ 1.
- **KEYTRUDA** as monotherapy is indicated for the treatment of recurrent or metastatic head and neck squamous cell carcinoma in adults whose tumours express PD-L1 with a ≥ 50% TPS and progressing on or after platinum-containing chemotherapy.

Renal cell carcinoma (RCC)

- KEYTRUDA, in combination with axitinib, is indicated for the first-line treatment of advanced renal cell carcinoma in adults.
- KEYTRUDA, in combination with lenvatinib, is indicated for the first-line treatment of advanced renal cell carcinoma in adults.
- **KEYTRUDA** as monotherapy is indicated for the adjuvant treatment of adults with renal cell carcinoma at increased risk of recurrence following nephrectomy or following nephrectomy and resection of metastatic lesions.

Microsatellite instability high (MSI-H) or mismatch repair deficient (dMMR) cancers

Colorectal cancer (CRC)

- **KEYTRUDA** as monotherapy is indicated for adults with MSI-H or dMMR colorectal cancer in the following settings:
 - first-line treatment of metastatic colorectal cancer;
 - treatment of unresectable or metastatic colorectal cancer after previous fluoropyrimidine-based combination therapy

Non-colorectal cancers

- **KEYTRUDA** as monotherapy is indicated for the treatment of the following MSI-H or dMMR tumours in adults with:
 - advanced or recurrent endometrial carcinoma, who have disease progression on or following prior treatment with a platinum-containing therapy in any setting and who are not candidates for curative surgery or radiation;
 - unresectable or metastatic gastric, small intestine, or biliary cancer, who have disease progression on or following at least one prior therapy.





KEYTRUDA[®] (pembrolizumab) indications¹

Oesophageal carcinoma

• **KEYTRUDA**, in combination with platinum and fluoropyrimidine-based chemotherapy, is indicated for the first-line treatment of locally advanced unresectable or metastatic carcinoma of the oesophagus in adults whose tumours express PD-L1 with a CPS ≥ 10.

Triple-negative breast cancer (TNBC)

- **KEYTRUDA**, in combination with chemotherapy as neoadjuvant treatment, and then continued as monotherapy as adjuvant treatment after surgery, is indicated for the treatment of adults with locally advanced, or early-stage triple-negative breast cancer at high risk of recurrence.
- **KEYTRUDA**, in combination with chemotherapy, is indicated for the treatment of locally recurrent unresectable or metastatic triple-negative breast cancer in adults whose tumours express PD-L1 with a CPS ≥ 10 and who have not received prior chemotherapy for metastatic disease.

Endometrial carcinoma (EC)

- KEYTRUDA, in combination with carboplatin and paclitaxel, is indicated for the first-line treatment of primary advanced or recurrent endometrial carcinoma in adults.
- **KEYTRUDA**, in combination with lenvatinib, is indicated for the treatment of advanced or recurrent endometrial carcinoma in adults who have disease progression on or following prior treatment with a platinum-containing therapy in any setting and who are not candidates for curative surgery or radiation.

Cervical cancer (CC)

- **KEYTRUDA**, in combination with chemoradiotherapy (external beam radiation therapy followed by brachytherapy), is indicated for the treatment of FIGO 2014 Stage III-IVA locally advanced cervical cancer in adults who have not received prior definitive therapy.
- **KEYTRUDA**, in combination with chemotherapy with or without bevacizumab, is indicated for the treatment of persistent, recurrent, or metastatic cervical cancer in adults whose tumours express PD-L1 with a CPS ≥ 1.

Gastric or gastro-oesophageal junction (GEJ) adenocarcinoma

- **KEYTRUDA**, in combination with trastuzumab, fluoropyrimidine and platinum-containing chemotherapy, is indicated for the first-line treatment of locally advanced unresectable or metastatic HER2-positive gastric or gastro-oesophageal junction adenocarcinoma in adults whose tumours express PD-L1 with a CPS ≥ 1.
- **KEYTRUDA**, in combination with fluoropyrimidine and platinum-containing chemotherapy, is indicated for the first-line treatment of locally advanced unresectable or metastatic HER2negative gastric or gastro-oesophageal junction adenocarcinoma in adults whose tumours express PD-L1 with a CPS ≥ 1

Biliary tract carcinoma (BTC)

• KEYTRUDA, in combination with gemcitabine and cisplatin, is indicated for the first-line treatment of locally advanced unresectable or metastatic biliary tract carcinoma in adults.









Mechanism of Action

KEYTRUDA Monotherapy Mechanism of Action

KEYTRUDA and Chemotherapy Mechanism of Action

KEYTRUDA + TKI Mechanism of Action



KEYTRUDA immune-mediated Adverse Events



KEYTRUDA immune-mediated Adverse Events Summary





Mechanism of Action







KEYTRUDA monotherapy mechanism of action^{1,2}

 KEYTRUDA is a humanised monoclonal antibody which binds to the programmed cell death-1 (PD-1) receptor, blocking both immune-suppressing ligands, PD-L1 and PD-L2, from interacting with PD-1 to help potentiate T-cell response and immune response^{1,2}





Images adapted from Harvey, RD (2014)



T cell reactivation with KEYTRUDA

KEYTRUDA binds to the PD-1 receptor and blocks its interaction with PD-L1 and PD-L2, which helps potentiate the immune response. While having an effect on the tumour, this could also affect the immune system's interaction with normal healthy cells throughout the body.

Normal immune response

When interacting with most aberrant or cancerous cells, T cells are activated and can attack tumour cells.

Tumour evasion and T cell deactivation

A proportion of tumours can evade the immune system through the PD-1 pathway. The PD-L1 and PD-L2 ligands on tumours can bind with PD-1 receptors on T cells to inactivate the T cells.

KEYTRUDA and cytotoxic chemotherapy mechanism of action^{3,4}

Cytotoxic chemotherapy can increase the rate of cancer cell death, activating a broad range of T-cells.

Combining **KEYTRUDA** and cytotoxic chemotherapy may enhance the immune response.



KEYTRUDA and TKI mechanisms of action^{1,5}

KEYTRUDA and TKIs, such as **axitinib**, inhibit two distinct disease pathways in advanced RCC.^{1,5}

KEYTRUDA

Axitinib

Tumour evasion of immune responses¹

KEYTRUDA binds to the PD-1 receptor potentiating T-cell responses, including anti-tumour response.

Tumour angiogenesis⁵

Axitinib is a potent inhibitor of VEGFR-1/2/3, receptors that have been implicated in pathologic angiogenesis, tumour growth, and metastatic progression of cancer.





KEYTRUDA and multi-TKI mechanisms of action¹

KEYTRUDA and multi-TKIs, such as **lenvatinib**, combine to help overcome resistance to immunotherapy.¹

KEYTRUDA

Tumour evasion of immune responses¹

KEYTRUDA binds to the PD-1 receptor potentiating T-cell responses, including anti-tumour response.

Lenvatinib

Greater T-cell activation

Lenvatinib in combination with KEYTRUDA results in a tumour microenvironment with greater T-cell activation to help overcome primary and acquired resistance to immunotherapy. This may improve tumour responses compared to either treatment alone.¹





Immune-Mediated Adverse Events (imAEs)

KEYTRUDA is associated with imAEs and these slides focus on the treatment modifications for imAEs only. A full list of undesirable effects can be found in the KEYTRUDA SmPC.

Consider all adverse events (AEs) and refer to the KEYTRUDA SmPC and Risk Minimisation Materials (RMM) for full information on all AEs before prescribing KEYTRUDA.







KEYTRUDA safety in a studied population¹

KEYTRUDA safety data studied populations based on Q3W administration



Pooled safety population of KEYTRUDA¹

- The safety of pembrolizumab as monotherapy has been evaluated in 7,631 patients across tumour types and across four doses (2 mg/kg bw every 3 weeks, 200 mg every 3 weeks, or 10 mg/kg bw every 2 or 3 weeks) in clinical studies
- types and across four doses (2 mg/kg bw every 3 weeks, 200 mg every 3 weeks, or 10 mg/kg bw every 2 or 3 weeks) in clinical studies.
- The safety of KEYTRUDA in combination with chemotherapy or chemoradiotherapy (CRT) has been evaluated in 6,093 patients across tumour types receiving 200 mg, 2 mg/kg bw or 10 mg/kg bw KEYTRUDA every 3 weeks, in clinical studies
- When KEYTRUDA is administered in combination with chemotherapy, refer to the SmPC for the respective combination therapy components prior to initiation of treatment





KEYTRUDA safety in a studied population¹

KEYTRUDA safety data studied populations based on Q3W administration



KEYTRUDA in combination with TKIs¹

- The safety of KEYTRUDA in combination with axitinib or lenvatinib in advanced RCC, and in combination with lenvatinib in advanced
 EC has been evaluated in a total of 1,456 patients with advanced RCC or advanced EC receiving 200 mg KEYTRUDA every
 3 weeks with either axitinib 5mg twice daily or lenvatinib 20 mg once daily in clinical studies, as appropriate
- When pembrolizumab is administered in combination with axitinib or lenvatinib, refer to the SmPC for axitinib or lenvatinib prior to initiation of treatment. For additional lenvatinib safety information related to advanced RCC see the SmPC for Kisplyx and for advanced EC see the SmPC for Lenvima





Urothelial carcinoma - specific precaution¹



Use of KEYTRUDA in urothelial carcinoma patients who have received prior platinum-containing chemotherapy¹

 Physicians should consider the delayed onset of KEYTRUDA effect before initiating treatment in patients with poorer prognostic features and/or aggressive disease. In urothelial carcinoma, a higher number of deaths within 2 months was observed in KEYTRUDA compared to chemotherapy. Factors associated with early deaths were fast progressive disease on prior platinum therapy and liver metastases





KEYTRUDA safety in a studied population¹

Certain groups were excluded from clinical trials



Patients with the following conditions were excluded from clinical studies¹:

- Active CNS metastases; ECOG PS ≥2 (except for urothelial carcinoma and RCC); HIV infection, hepatitis B or hepatitis C infection (except for BTC); active systemic autoimmune disease; interstitial lung disease; prior pneumonitis requiring systemic corticosteroid therapy; a history of severe hypersensitivity to another monoclonal antibody; receiving immunosuppressive therapy and a history of severe immune-mediated adverse reactions from treatment with ipilimumab, defined as any Grade 4 toxicity or Grade 3 toxicity requiring corticosteroid treatment (>10 mg/day prednisone or equivalent) for greater than 12 weeks
- Patients with active infections were excluded from clinical studies and were required to have their infection treated prior to receiving KEYTRUDA. Patients with active infections occurring during treatment with KEYTRUDA were managed with appropriate medical therapy. Patients with clinically significant renal (creatinine >1.5 x ULN) or hepatic (bilirubin >1.5 x ULN, ALT, AST >2.5 x ULN in the absence of liver metastases) abnormalities at baseline were excluded from clinical studies, therefore information is limited in patients with severe renal and moderate to severe hepatic impairment





KEYTRUDA immune-mediated adverse event overview¹



KEYTRUDA is associated with immune-mediated adverse events.¹

Recommended treatment modifications for KEYTRUDA are dependent on type and Grade. Specific immune-mediated adverse events can be found in the SmPC and later in this document.

- Immune-mediated adverse reactions, including severe and fatal cases, have occurred in patients receiving KEYTRUDA
- Most immune-mediated adverse events are reversible and may be managed with interruptions of KEYTRUDA, administration of corticosteroids and/or supportive care
- Risk of immune-mediated adverse reactions following immune-checkpoint inhibitor therapy may be increased in patients with preexisting autoimmune disease (AID).
- Immune-mediated adverse events have also occurred after the last dose of KEYTRUDA. Immune-mediated adverse reactions affecting more than one body system can occur simultaneously



KEYTRUDA Grade 3–4 immune-mediated adverse events.^{1,6}

Toxicity grades are in accordance with National Cancer Institute Common Terminology Criteria for Adverse Events Version 4.0 (NCI-CTCAE v.4). A table of the grades for individual immune-mediated adverse events can be found <u>HERE</u> and at the end of this presentation.

- KEYTRUDA must be permanently discontinued for any Grade 3 immune-mediated adverse reaction that recurs and for any Grade 4 immune-mediated adverse reaction toxicity, except for endocrinopathies that are controlled with replacement hormones
- For more details, please refer to the SmPC for managing individual immune-mediated adverse reactions

Please refer to the KEYTRUDA Summary of Product Characteristics and Risk Minimisation Materials before prescribing KEYTRUDA.





Treatment Modifications for Immune-Mediated Adverse Events (imAEs)^{1,6}





Please refer to the KEYTRUDA SmPC and Risk Minimisation Materials (RMM) for full information on managing AEs before prescribing KEYTRUDA.





Hyperthyroidism









Hyperthyroidism¹



Treatment modifications for hyperthyroidism in patients prescribed KEYTRUDA

		Grade 1	Grade 2	Grade 3–4
*	KEYTRUDA treatment	May continue treatment and monitor.	May continue treatment and monitor.	Withhold until recovery to Grade ≤1. If toxicity does not resolve to Grade 0–1 within 12 weeks after last dose of KEYTRUDA, or corticosteroid dosing cannot be reduced to ≤10 mg prednisone or equivalent per day within 12 weeks, permanently discontinue KEYTRUDA.
	Corticosteroid treatment	-	-	Consider administration of corticosteroids and taper as required. Please refer to the SmPC for further information.
•	Additional management	May be managed symptom Thyroid function and hormo monitored to ensure appro	natically. one levels should be priate replacement.	For patients with Grade 3 or Grade 4 endocrinopathies that improved to Grade 2 or lower and are controlled with hormone replacement, if indicated, continuation of KEYTRUDA may be considered after corticosteroid taper, if needed. Otherwise treatment should be discontinued.

Long-term hormone replacement therapy may be necessary in cases of immune-mediated endocrinopathies.



Hyperthyroidism¹



Prevalence in clinical trials



Pooled safety population of KEYTRUDA¹

Hyperthyroidism occurred in 394 (5.2%) patients, including Grade 2 or 3 cases in 108 (1.4%) and 9 (0.1% patients, respectively, receiving pembrolizumab. The median time to onset of hyperthyroidism was 1.4 months (range 1 day to 23.2 months). The median duration was 1.6 months (range 4 days to 43.1+ months). Hyperthyroidism led to discontinuation of pembrolizumab in 4 (0.1%) patients. Hyperthyroidism resolved in 326 (82.7%) patients, 11 with sequelae)



Monitoring hyperthyroidism in patients prescribed KEYTRUDA¹

- Patients should be monitored for changes in thyroid function (at the start of treatment, periodically throughout and when indicated by clinical evaluation) as well as clinical signs and symptoms of thyroid disorders. Hormone levels should also be monitored
- Along with hypothyroidism/hyperthyroidism, thyroiditis has also been reported and can occur at any time during treatment





Hypothyroidism









Hypothyroidism¹



Treatment modifications for hypothyroidism in patients prescribed KEYTRUDA

KEYTRUDA treatment	May continue treatment and monitor
Additional management	May be managed with replacement hormone therapy without treatment interruption and corticosteroids Thyroid function and hormone levels should be monitored to ensure appropriate hormone replacement

Long-term hormone replacement therapy may be necessary in cases of immune-mediated endocrinopathies.





Hypothyroidism¹



Prevalence in clinical trials



Pooled safety population of KEYTRUDA¹

- Hypothyroidism occurred in 939 (12.3%) patients, including Grade 2 or 3 cases in 687 (9.0%) and 8 (0.1%) patients, respectively, receiving pembrolizumab. The median time to onset of hypothyroidism was 3.4 months (range 1 day to 25.9 months). The median duration was not reached (range 2 days to 63.0+ months). Hypothyroidism led to discontinuation of pembrolizumab in 6 (0.1%) patients. Hypothyroidism resolved in 216 (23.0%) patients, 16 with sequelae.
- In patients with cHL (n=389) the incidence of hypothyroidism was 17%, all of which were Grade 1 or 2. In patients with HNSCC treated with pembrolizumab as monotherapy (n=909), the incidence of hypothyroidism was 16.1% (all Grades) with 0.3% Grade 3. In patients with HNSCC treated with pembrolizumab in combination with platinum and 5-FU chemotherapy (n=276), the incidence of hypothyroidism was 15.2%, all of which were Grade 1 or 2. In patients treated with pembrolizumab in combination or lenvatinib (n=1,456), the incidence of hypothyroidism was 46.2% (all Grades) with 0.8% Grade 3 or 4.



Monitoring hypothyroidism in patients prescribed KEYTRUDA¹

- Patients should be monitored for the changes in thyroid function (at the start of treatment, periodically throughout and when indicated by clinical evaluation) as well as clinical signs and symptoms of thyroid disorders. Hormone levels should also be monitored
- Along with hypothyroidism/hyperthyroidism, thyroiditis has also been reported and can occur at any time during treatment
- Hypothyroidism is more frequently reported in patients with HNSCC with prior radiation therapy













• -----



UK PI

88

Treatment modifications for colitis in patients prescribed KEYTRUDA

	Grade 1	Grade 2–3	Grade 4
KEYTRUDA treatment	May continue treatment and monitor.	Withhold until adverse event recovers to Grade 0–1.	Permanently discontinue.
		If toxicity does not resolve to Grade 0–1 within 12 weeks after last dose of KEYTRUDA, or corticosteroid dosing cannot be reduced to ≤10 mg prednisone or equivalent per day within 12 weeks, permanently discontinue KEYTRUDA. For recurrent Grade 3, permanently discontinue.	
Corticosteroid treatment	-	Initial dose of 1–2 mg/kg per day predr taper.	isone or equivalent followed by a
Additional management	The potential risk of gastrointestinal pe	rforation should be taken into considerat	ion.





Prevalence in clinical trials



Pooled safety population of KEYTRUDA¹

- Colitis occurred in 158 (2.1%) patients, including Grade 2, 3 or 4 cases in 49 (0.6%), 82 (1.1%) and 6 (0.1%) patients, respectively, receiving pembrolizumab. The median time to onset of colitis was 4.3 months (range 2 days to 24.3 months). The median duration was 1.1 month (range 1 day to 45.2 months). Colitis led to discontinuation of pembrolizumab in 48 (0.6%) patients. Colitis resolved in 132 patients, 2 with sequelae.
- In patients with CRC treated with KEYTRUDA as monotherapy (n=153), the incidence of colitis was 6.5% (all Grades) with 2.0%
 Grade 3 and 1.3% Grade 4.



Monitoring colitis in patients prescribed KEYTRUDA¹

Patients should be monitored for the signs and symptoms of colitis and other causes excluded.





Hepatitis













Treatment modifications for hepatitis in patients prescribed KEYTRUDA

		Grade 1	Grade 2 (AST or ALT >3x to 5x ULN or total bilirubin >1.5x to 3x ULN)	Grade 3–4 (AST or ALT >5x ULN or total bilirubin >3x ULN)
* -	KEYTRUDA treatment	May continue treatment and monitor.	Withhold until adverse event recovers to Grade 0–1. If toxicity does not resolve to Grade 0–1 within 12 weeks after last dose of KEYTRUDA, or corticosteroid dosing cannot be reduced to ≤10 mg prednisone or equivalent per day within 12 weeks, permanently discontinue KEYTRUDA.	Permanently discontinue.
	Corticosteroid treatment	-	Initial dose of 0.5–1 mg/kg per day prednisone or equivalent followed by a taper.	Dose of 1–2 mg/kg per day prednisone or equivalent followed by a taper.
	Additional management	In the case of liver meta and last ≥1 week, perm	stasis with baseline Grade 2 elevation of AST or AL anently discontinue KEYTRUDA.	T, hepatitis with AST or ALT increases ≥50%









Prevalence in clinical trials



Patients with clinically significant renal (creatinine >1.5 x ULN) or hepatic (bilirubin >1.5 x ULN, ALT, AST >2.5 x ULN in the absence of liver metastases) abnormalities at baseline were excluded from clinical trials, therefore information is limited in patients with severe renal and moderate to severe hepatic impairment.



Pooled safety population of KEYTRUDA¹

Hepatitis occurred in 80 (1.0%) patients, including Grade 2, 3 or 4 cases in 12 (0.2%), 55 (0.7%) and 8 (0.1%) patients, respectively, receiving pembrolizumab. The median time to onset of hepatitis was 3.5 months (range 8 days to 26.3 months). The median duration was 1.3 months (range 1 day to 29.0+ months). Hepatitis led to discontinuation of pembrolizumab in 37 (0.45%) patients. Hepatitis resolved in 60 patients



Monitoring hepatitis in patients prescribed KEYTRUDA¹

- Patients should be monitored for changes in liver function (at the start of treatment, periodically throughout and when indicated by clinical evaluation) and for symptoms of hepatitis. When hepatitis is suspected, exclude other causes of liver dysfunction
- Please refer to the KEYTRUDA SmPC for full information on managing elevation in liver enzymes before prescribing KEYTRUDA





Liver enzyme elevations









Liver enzyme elevations¹



Treatment modifications for liver enzyme elevations in patients with RCC prescribed KEYTRUDA in combination with axitinib

		AST or ALT <3x ULN	AST or ALT ≥3x ULN but <10x ULN without concurrent total bilirubin ≥2x ULN	AST or ALT >5 x ULN or total bilirubin >3 x ULN
Ŷ	KEYTRUDA treatment	May continue treatment and monitor enzymes as long as below limits detailed here.	Withhold until adverse event recovers to Grade 0–1.	Permanently discontinue.
	Corticosteroid treatment	-	Consider administration of corticosteroids and taper as required.	Consider administration of corticosteroids and taper as required.
•	Additional management	If treatment is withheld f ULN, rechallenge with a If rechallenging with axi	following AST or ALT ≥3 times ULN but <10 times a single medicine or sequential rechallenge with be tinib, consider dose reduction as per axitinib SmP	ULN without concurrent total bilirubin ≥2 times oth medicines after recovery may be considered. C.
		If ALT or AST ≥ 10 times axitinib should be perma	s ULN or > 3 times ULN with concurrent total biliru anently discontinued and corticosteroid therapy m	lbin ≥ 2 times ULN, both KEYTRUDA and ay be considered.

Liver enzyme elevations¹



Prevalence in clinical trials



Patients with clinically significant renal (creatinine >1.5 x ULN) or hepatic (bilirubin >1.5 x ULN, ALT, AST >2.5 x ULN in the absence of liver metastases) abnormalities at baseline were excluded from clinical trials, therefore information is limited in patients with severe renal and moderate to severe hepatic impairment.



Pooled safety population of KEYTRUDA¹

In a clinical study of previously untreated patients with RCC receiving pembrolizumab in combination with axitinib, a higher than expected incidence of Grades 3 and 4 ALT increased (20%) and AST increased (13%) were observed. The median time to onset of ALT increased was 2.3 months (range: 7 days to 19.8 months). In patients with ALT \geq 3 times ULN (Grades 2–4, n=116), ALT resolved to Grades 0–1 in 94%. Fifty-nine percent of the patients with increased ALT received systemic corticosteroids. Of the patients who recovered, 92 (84%) were rechallenged with either pembrolizumab (3%) or axitinib (31%) monotherapy or with both (50%). Of these patients, 55% had no recurrence of ALT >3 times ULN, and of those patients with recurrence of ALT >3 times ULN, all recovered. There were no Grade 5 hepatic events.



Monitoring liver enzyme elevations in patients prescribed KEYTRUDA¹

- Patients should be monitored for changes in liver enzymes before initiation of and periodically throughout treatment. More frequent monitoring of liver enzymes may be considered when not used as a monotherapy.
- Please refer to the KEYTRUDA SmPC for full information on managing elevation in liver enzymes before prescribing KEYTRUDA.





Adrenal insufficiency









Adrenal insufficiency¹



Treatment modifications for adrenal insufficiency in patients prescribed KEYTRUDA

		Grade 1	Grade 2	Grade 3–4
	KEYTRUDA	May continue	Withhold treatment until	Withhold until recovery to Grade ≤1.
	treatment	treatment and monitor.	controlled by hormone replacement.	If toxicity does not resolve to Grade 0–1 within 12 weeks after last dose of KEYTRUDA, or corticosteroid dosing cannot be reduced to ≤10 mg prednisone or equivalent per day within 12 weeks, permanently discontinue KEYTRUDA.
	Corticosteroid treatment	Administer corticosteroi	minister corticosteroids to treat adrenal insufficiency, followed by corticosteroid taper if indicated.	
Additional management Hormone replacement therapy if i		herapy if indicated.	For patients with Grade 3 or Grade 4 endocrinopathies that improved to Grade 2 or lower and are controlled with hormone replacement, if indicated, continuation of KEYTRUDA may be considered after corticosteroid taper, if needed.	
				Otherwise treatment should be discontinued.

Long-term hormone replacement therapy may be necessary in cases of immune-mediated endocrinopathies.

Adrenal insufficiency¹



Prevalence in clinical trials



Pooled safety population of KEYTRUDA¹

Adrenal insufficiency occurred in 74 (1.0%) patients, including Grade 2, 3 or 4 cases in 34 (0.4%), 31 (0.4%) and 4 (0.1%) patients, respectively, receiving pembrolizumab. The median time to onset of adrenal insufficiency was 5.4 months (range 1 day to 23.7 months). The median duration was not reached (range 3 days to 40.1+ months). Adrenal insufficiency led to discontinuation of pembrolizumab in 13 (0.2%) patients. Adrenal insufficiency resolved in 28 patients, 11 with sequelae



Monitoring adrenal insufficiency in patients prescribed KEYTRUDA¹

- Patients should be monitored for the signs and symptoms of adrenal insufficiency and exclude other causes
- Pituitary function and hormone levels should be monitored to ensure appropriate hormone replacement





Hypophysitis









Hypophysitis¹



<u>UK</u>

ΡΙ

Treatment modifications for hypophysitis in patients prescribed KEYTRUDA

		Grade 1	Grade 2	Grade 3–4 or symptomatic hypophysitis
*	KEYTRUDA treatment	May continue treatment and monitor.	Withhold treatment until controlled by hormone replacement	Withhold until recovery to Grade ≤1. If toxicity does not resolve to Grade 0–1 within 12 weeks after last dose of KEYTRUDA, or corticosteroid dosing cannot be reduced to ≤10 mg prednisone or equivalent per day within 12 weeks, permanently discontinue KEYTRUDA.
	Corticosteroid treatment	Administer corticosteroi	ds to treat hypophysitis, followed by cort	icosteroid taper if needed.
•	Additional management	Hormone replacement therapy if indicated.		For patients with Grade 3 or Grade 4 endocrinopathies that improved to Grade 2 or lower and are controlled with hormone replacement, if indicated, continuation of KEYTRUDA may be considered after corticosteroid taper, if needed. Otherwise treatment should be discontinued.

Long-term hormone replacement therapy may be necessary in cases of immune-mediated endocrinopathies.

Hypophysitis¹



Prevalence in clinical trials



Pooled safety population of KEYTRUDA¹

Hypophysitis occurred in 52 (0.7%) patients, including Grade 2, 3 or 4 cases in 23 (0.3%), 24 (0.3%) and 1 (<0.1%) patients, respectively, receiving pembrolizumab. The median time to onset of hypophysitis was 5.9 months (range 1 day to 17.7 months). The median duration was 3.6 months (range 3 days to 48.1+ months). Hypophysitis led to discontinuation of pembrolizumab in 14 (0.2%) patients. Hypophysitis resolved in 23 patients, 8 with sequelae</p>



Monitoring hypophysitis in patients prescribed KEYTRUDA¹

- Patients should be monitored for the signs and symptoms of hypophysitis (including hypopituitarism) and exclude other causes
- Pituitary function and hormone levels should be monitored to ensure appropriate hormone replacement





Pneumonitis













Treatment modifications for pneumonitis in patients prescribed KEYTRUDA

	Grade 1	Grade 2	Grade 3–4
KEYTRUDA treatment	May continue treatment and monitor.	Withhold until adverse event recovers to Grade 0–1. If toxicity does not resolve to Grade 0–1 within 12 weeks after last dose of KEYTRUDA, or corticosteroid dosing cannot be reduced to ≤10 mg prednisone or equivalent per day within 12 weeks, permanently discontinue KEYTRUDA. For recurrent Grade 2, permanently discontinue.	Permanently discontinue.
Corticosteroid treatment	-	Dose of 1–2 mg/kg per day prednisone or equivalent for	ollowed by a taper.
Additional management	Note: fatal cases of pneumonitis have been reported in patients receiving KEYTRUDA.		

KEYTRUDA should be permanently discontinued for Grade 3–4 or recurrent Grade 2 pneumonitis





Pneumonitis¹



Prevalence in clinical trials



Pooled safety population of KEYTRUDA¹

- Pneumonitis occurred in 324 (4.42%) patients, including Grade 2, 3, 4 or 5 cases in 143 (1.9%), 81 (1.1%), 19 (0.2%) and 9 (0.1%) patients, respectively, receiving pembrolizumab. The median time to onset of pneumonitis was 3.7 months (range 2 days to 27.2 months). The median duration was 2.0 months (range 1 day to 51.0+ months). Pneumonitis occurred more frequently in patients with a history of prior thoracic radiation (8.1%) than in patients who did not receive prior thoracic radiation (3.9%). Pneumonitis led to discontinuation of pembrolizumab in 131 (1.7%) patients. Pneumonitis resolved in 196 patients, 6 with sequelae
- In patients with NSCLC, pneumonitis occurred in 230 (6.1%), including Grade 2, 3, 4 or 5 cases in 103 (2.7%), 63 (1.7%), 17 (0.4%) and 10 (0.3%), respectively. In patients with NSCLC, pneumonitis occurred in 8.9% with a history of prior thoracic radiation
- In patients with cHL, the incidence of pneumonitis (all Grades) ranged from 5.2% to 10.8% for cHL patients in KEYNOTE-087 (n=210) and KEYNOTE-204 (n=148), respectively



Monitoring pneumonitis in patients prescribed KEYTRUDA¹

Patients should be monitored for signs and symptoms of pneumonitis. When pneumonitis is suspected, evaluate with radiographic imaging to exclude other causes





Type 1 Diabetes Mellitus (Hyperglycaemia/Diabetic Ketoacidosis)









Type 1 diabetes mellitus¹ (Hyperglycaemia/Diabetic ketoacidosis)



Treatment modifications for type 1 diabetes in patients prescribed KEYTRUDA

		Type 1 diabetes	Type 1 diabetes associated with <u>Grade ≥3</u> hyperglycaemia (glucose >250 mg/dL or >13.9 mmol/L) or ketoacidosis
	KEYTRUDA treatment	May continue treatment and monitor.	KEYTRUDA should be withheld until metabolic control is achieved.
\frown			
	Additional management	Administer insulin.	Administer insulin.

Long-term hormone replacement therapy may be necessary in cases of immune-mediated endocrinopathies.





Type 1 diabetes mellitus¹ (Hyperglycaemia/Diabetic ketoacidosis)



Prevalence in clinical trials



Pooled safety population of KEYTRUDA¹

- Severe endocrinopathies, including type 1 diabetes mellitus, have been observed with KEYTRUDA treatment
- In patients treated with KEYTRUDA monotherapy, the proportion of patients who experienced a shift from baseline to a Grade 3 or 4 laboratory abnormality for increased glucose (potentially indicating hyperglycaemia) was 4.6% for KEYTRUDA monotherapy, 5.5% for KEYTRUDA in combination with chemotherapy or CRT and 7.8% for KEYTRUDA in combination with axitinib or lenvatinib



Monitoring hyperglycaemia in patients prescribed KEYTRUDA¹

Patients should be monitored for hyperglycaemia or other signs and symptoms of type 1 diabetes





Nephritis













Treatment modifications for nephritis in patients prescribed KEYTRUDA

\frown		Grade 1	Grade 2 (creatinine >1.5 to ≤3x ULN)	Grade 3–4 (creatinine ≥3 x ULN)
	KEYTRUDA treatment	May continue treatment and monitor.	Withhold based on the severity of creatinine elevations and until adverse events recover to Grade 0–1.	Permanently discontinue.
			If toxicity does not resolve to Grade 0–1 within 12 weeks after last dose of KEYTRUDA, or corticosteroid dosing cannot be reduced to ≤10 mg prednisone or equivalent per day within 12 weeks, permanently discontinue KEYTRUDA.	
	O anti a a sta nai d		Initial data of 1. O man/len man data and data	
	treatment	-	Initial dose of 1–2 mg/kg per day predhisone d	or equivalent followed by a taper.
\bigcirc				
	Additional management	Management depends on the sever	rity of creatinine elevations.	





Nephritis¹



Prevalence in clinical trials



Pooled safety population of KEYTRUDA¹

- Nephritis occurred in 37 (0.5%) patients, including Grade 2, 3 or 4 cases in 11 (0.1%), 19 (0.2%) and 2 (< 0.1%) patients, respectively, receiving pembrolizumab as monotherapy. The median time to onset of nephritis was 4.2 months (range 12 days to 21.4 months). The median duration was 3.3 months (range 6 days to 28.2+ months). Nephritis led to discontinuation of pembrolizumab in 17 (0.2%) patients. Nephritis resolved in 25 patients, 5 with sequelae.
- In patients with non-squamous NSCLC treated with pembrolizumab in combination with pemetrexed and platinum chemotherapy (n=488), the incidence of nephritis was 1.4% (all Grades) with 0.8% Grade 3 and 0.4% Grade 4



Monitoring nephritis in patients prescribed KEYTRUDA¹

Patients should be monitored for changes in renal function. When nephritis is suspected, exclude other causes of renal dysfunction





Other Immune-Mediated Adverse Events (imAEs)









Other immune-mediated adverse events¹



Treatment modifications for other imAEs in patients prescribed KEYTRUDA

	Grade 2–3*	Grade 4 and recurrent Grade 3
 KEYTRUDA treatment 	 Based on the severity and type of the reaction, withhold treatment until adverse reactions recover to Grade 0–1. 	Permanently discontinue
	• Permanently discontinue if a Grade 3 event occurs more than once	
	 First occurrence of Grade 3 myocarditis or encephalitis or Guillain-Barré syndrome – permanently discontinue 	
 Corticosteroid treatment 	Administer corticosteroids followed by corticosteroid t	aper if indicated.

D	Add	ditional management	*In the case of Grade 2 and first occurrence of Grade 3 adverse events: If treatment related toxicity is not resolved within 12 weeks after last dose of KEYTRUDA or if corticosteroid dose cannot be reduced less or equal to 10 mg/day prednisolone or equivalent per day within 12 weeks, KEYTRUDA should be permanently discontinued
			RETTRODA Should be permanently discontinued.





Other immune-mediated adverse events¹



(U)

Monitoring other immune-mediated adverse events in patients prescribed KEYTRUDA¹

- Patients should be monitored for the signs and symptoms of other immune-mediated adverse events. Other causes should be excluded
- The following additional clinically significant, immune-mediated adverse reactions have been reported in clinical studies or in postmarketing experience: uveitis, arthritis, myositis, myocarditis, pancreatitis, Guillain-Barré syndrome, myasthenic syndrome, haemolytic anaemia, sarcoidosis, encephalitis, myelitis, vasculitis, cholangitis sclerosing, gastritis, cystitis noninfective and hypoparathyroidism and pericarditis
- Immune-mediated adverse reactions affecting more than one body system can occur simultaneously
- Severe and fatal cases have been reported in clinical trials or in post-marketing experience





Transplant-related adverse reactions¹

Solid organ transplant rejection¹

- Solid organ transplant rejection has been reported in the post-marketing setting in patients treated with PD-1 inhibitors. Treatment with KEYTRUDA may increase the risk of rejection in solid organ transplant recipients.
- The benefit of treatment with KEYTRUDA vs the risk of possible organ rejection should be considered in these patients.





Transplant-related adverse reactions¹

Complications of allogeneic haematopoietic stem cell transplant (HSCT)¹

Allogeneic HSCT after treatment with KEYTRUDA

- Cases of graft-versus-host-disease (GVHD) and hepatic veno-occlusive disease (VOD) have been observed in patients with cHL undergoing allogeneic HSCT after previous exposure to pembrolizumab
- Until further data become available, careful consideration to the potential benefits of HSCT and the possible increased risk of transplantrelated complications should be made case by case

Allogeneic HSCT prior to treatment with KEYTRUDA

- In patients with a history of allogeneic HSCT, acute GVHD, including fatal GVHD, has been reported after treatment with pembrolizumab.
 Patients who experienced GVHD after their transplant procedure may be at an increased risk for GVHD after treatment with pembrolizumab
- · Consider the benefit of treatment with pembrolizumab versus the risk of possible GVHD in patients with a history of allogeneic HSCT

Allogeneic HSCT in classical Hodgkin lymphoma (cHL)

- Of 14 patients in KEYNOTE-013 who proceeded to allogeneic HSCT after treatment with pembrolizumab, 6 patients reported GVHD and 1 patient reported chronic GVHD, none of which were fatal. Two patients experienced hepatic VOD, one of which was fatal. One patient experienced engraftment syndrome post-transplant
- Of 32 patients in KEYNOTE-087 who proceeded to allogeneic HSCT after treatment with pembrolizumab, 16 patients reported acute GVHD and 7 patients reported chronic GVHD, two of which were fatal. No patients experienced hepatic VOD. No patients experienced engraftment syndrome post-transplant
- Of 14 patients in KEYNOTE-204 who proceeded to allogeneic HSCT after treatment with pembrolizumab, 8 patients reported acute GVHD and 3 patients reported chronic GVHD, none of which were fatal. No patients experienced hepatic VOD. One patient experienced engraftment syndrome post-transplant





Infusion-Related Reactions









Infusion-related reactions¹



Treatment modifications for infusion-related reactions in patients prescribed KEYTRUDA

		Mild to moderate reactions (Grade 1–2)	Severe reactions (Grade 3–4)
)	KEYTRUDA treatment	May continue treatment and monitor.	Stop infusion. Permanently discontinue.
	Additional management	Monitor closely and consider premedication with antipyretic and antihistamine therapy.	-

KEYTRUDA, when used as monotherapy or in combination, must be administered by intravenous infusion over 30 minutes.





Infusion-related reactions¹



Prevalence in clinical trials



Pooled safety population of KEYTRUDA¹

Severe infusion-related reactions, including hypersensitivity and anaphylaxis, have been reported in patients receiving KEYTRUDA.



Monitoring infusion-related reactions in patients prescribed KEYTRUDA¹

- Severe infusion-related reactions including hypersensitivity and anaphylaxis have been reported with patients receiving KEYTRUDA. These included drug hypersensitivity, anaphylactic reaction, anaphylactoid reaction, hypersensitivity and cytokine release syndrome
- Patients should be monitored during infusion





Use of KEYTRUDA in combination with chemotherapy¹





KEYTRUDA in combination with chemotherapy should be used with caution in patients ≥75 years after careful consideration of the potential benefit/risk on an individual basis





Skin Reactions











Treatment modifications for skin reactions in patients prescribed KEYTRUDA

	Grade 1–2	Grade 3 or <u>suspected</u> SJS or TEN	Grade 4 or <u>confirmed</u> SJS or TEN
KEYTRUDA treatment	May continue treatment and monitor.	Withhold until adverse reactions recover to Grades 0–1. If toxicity does not resolve to Grade 0–1 within 12 weeks after last dose of KEYTRUDA, or corticosteroid dosing cannot be reduced to ≤10 mg prednisone or equivalent per day within 12 weeks, permanently discontinue KEYTRUDA.	Permanently discontinue.



Additional	For signs or symptoms of SJS or TEN, the patient should be referred to a
management	specialised unit for assessment and treatment.





Skin reactions¹



Prevalence in clinical trials



Pooled safety population of KEYTRUDA¹

- Immune-mediated severe skin reactions occurred in 130 (1.7%) patients, including Grade 2, 3, 4 or 5 cases in 11 (0.1%), 103 (1.3%), 1 (<0.1%) and 1 (<0.1%) patients, respectively, receiving pembrolizumab. The median time to onset of severe skin reactions was 2.8 months (range 2 days to 25.5 months). The median duration was 1.9 months (range 1 day to 47.1+ months). Severe skin reactions led to discontinuation of pembrolizumab in 18 (0.2%) patients. Severe skin reactions resolved in 95 patients, 2 with sequelae
- Rare cases of SJS and TEN, some of them with fatal outcome, have been reported in patients receiving pembrolizumab
- The safety of pembrolizumab in combination with enfortumab vedotin has been evaluated among 564 patients with unresectable or metastatic urothelial carcinoma receiving 200 mg pembrolizumab on Day 1 and enfortumab vedotin 1.25 mg/kg on Days 1 and 8 of each 21-day cycle. The incidence of rash maculo-papular was 36% all Grades (10% Grades 3-4), which is higher than observed in pembrolizumab monotherapy.



Monitoring skin reactions in patients prescribed KEYTRUDA¹

- Patients should be monitored for suspected skin reactions. Other causes should be excluded
- Caution should be used when considering the use of KEYTRUDA in a patient who has previously experienced a severe or lifethreatening skin adverse reaction on prior treatment with other immune-stimulatory anticancer agents





Immune-Mediated Adverse Events Summary







Immune-mediated Adverse Event Summary¹

Hyperthyroidism	 For changes in thyroid function (at the start of treatment, periodically during treatment, and as indicated based on clinical evaluation) and clinical signs and symptoms of thyroid disorders 		 For Grade 2 hyperthyroidism, treatment with KEYTRUDA may continue with monitoring. For patients with Grade 3 or Grade 4 hyperthyroidism that improved to Grade 2 or lower, continuation may be considered, after corticosteroid taper, if needed. Thyroid function and hormone levels should be monitored to ensure appropriate hormone replacement For Grade ≥3 withhold until recovery to Grade ≤1.* For patients with Grade 3–4 endocrinopathies that improved to Grade 2 or lower and is controlled with hormone replacement, if indicated, continuation of KEYTRUDA may be considered after corticosteroid taper, if needed
Hypothyroidism	For changes in thyroid function		 Symptoms may be managed with replacement hormone therapy and treatment with KEYTRUDA may continue with monitoring Thyroid function and hormone levels should be monitored to ensure appropriate replacement
Immune- mediated colitis	 For signs and symptoms of colitis; exclude other causes 		 Withhold KEYTRUDA for Grade 2 or Grade 3 colitis until adverse reactions recover to Grades 0–1* Administer corticosteroids for Grade ≥2 events (initial dose of 1–2 mg/kg per day prednisone or equivalent followed by a taper) Permanently discontinue KEYTRUDA for recurrent Grade 3 or Grade 4 colitis
Immune- mediated hepatitis	• For changes in liver function (at the start of treatment, periodically during treatment, and as indicated based on clinical evaluation) and symptoms of hepatitis; exclude other causes	-	 For Grade 2 hepatitis, withhold until adverse event recovers to Grade 0–1.* Administer an initial dose of 0.5–1 mg/kg per day prednisone or equivalent followed by a taper For Grade ≥3 hepatitis, permanently discontinue KEYTRUDA. Administer 1–2 mg/kg per day prednisone or equivalent followed by a taper

*If treatment-related toxicity does not resolve to Grades 0–1 within 12 weeks after last dose of KEYTRUDA, or if corticosteroid dosing cannot be reduced to ≤10 mg prednisone or equivalent per day within 12 weeks, KEYTRUDA should be permanently discontinued.



Immune-mediated Adverse Event Summary¹

3,0)- SP)-	Immune-mediated endocrinopathies: adrenal insufficiency and hypophysitis	 For signs and symptoms of adrenal insufficiency and hypophysitis (including hypopituitarism) and other causes excluded 	 For Grade 2 adrenal insufficiency, and hypophysitis, withhold KEYTRUDA until controlled by hormone replacement For Grade ≥3 adrenal insufficiency and hypophysitis withhold KEYTRUDA until adverse reactions recover to Grades 0–1* For patients with Grade 3 or Grade 4 endocrinopathies that improved to Grade 2 or lower and are controlled with hormone replacement, if indicated, continuation of KEYTRUDA may be considered after corticosteroid taper, if needed. Otherwise treatment should be discontinued.
	Type 1 diabetes	 For hyperglycaemia or other signs and symptoms of diabetes Insulin should be administered for type 1 diabetes 	 For type 1 diabetes, KEYTRUDA may continue with insulin* For type 1 diabetes associated with Grade ≥3 hyperglycaemia or associated ketoacidosis, withhold KEYTRUDA. Treatment may be restarted if metabolic control is achieved
	Immune-mediated pneumonitis	 For signs and symptoms of pneumonitis Suspected pneumonitis should be confirmed with radiographic imaging and other causes excluded 	 Administer corticosteroids for Grade ≥2 events (initial dose of 1–2 mg/kg per day prednisone or equivalent followed by a taper) Withhold KEYTRUDA for Grade 2 pneumonitis until adverse reactions recover to Grades 0–1.* Permanently discontinue KEYTRUDA for Grade 3, Grade 4, or recurrent Grade 2 pneumonitis
	Immune-mediated nephritis	 For changes in renal function; exclude other causes 	 Administer corticosteroids for Grade ≥2 events (initial dose of 1–2 mg/kg per day prednisone or equivalent followed by a taper) Based on severity of creatinine elevations: Withhold KEYTRUDA for Grade 2 with creatinine >1.5 to ≤3 times upper limit of normal (ULN) until adverse reactions recover to Grades 0–1.* Permanently discontinue KEYTRUDA for Grade ≥ 3 nephritis with creatinine > 3 times ULN

*If treatment-related toxicity does not resolve to Grades 0–1 within 12 weeks after last dose of KEYTRUDA, or if corticosteroid dosing cannot be reduced to ≤10 mg prednisone or equivalent per day within 12 weeks, KEYTRUDA should be permanently discontinued.





Immune-mediated Adverse Event Summary¹

	Other immune- mediated adverse reactions	 Monitor for signs and symptoms of: uveitis, arthritis, myositis, myocarditis, pancreatitis, Guillain-Barré syndrome, myasthenic syndrome, haemolytic anaemia, sarcoidosis, encephalitis, myelitis, vasculitis, cholangitis sclerosing, gastritis, cystitis noninfective, hypoparathyroidism, and solid organ transplant rejection following KEYTRUDA treatment in donor organ recipients 	 For Grade 2–3, withhold KEYTRUDA until adverse reactions recover to Grade 0–1* KEYTRUDA should be permanently discontinued in the case of Grade 3 or 4 myocarditis, encephalitis and Guillain Barré syndrome Permanently discontinue drug if any Grade 3 immune-mediated toxicity occurs a second time and for any Grade 4 immune-mediated toxicity Based on limited data from clinical studies in patients whose immune-mediated adverse reactions could not be controlled with corticosteroid use, consider administration of other systemic immunosuppressants
	Infusion related reactions	 Severe infusion-related reactions, including hypersensitivity and anaphylaxis, have been reported in patients receiving KEYTRUDA 	 For Grade 1–2 (mild to moderate) infusion-related reactions, KEYTRUDA treatment may continue with monitoring. Monitor closely and consider premedication with antipyretic and antihistamine therapy For Grade 3–4 (severe) reactions, infusion should be stopped and KEYTRUDA permanently discontinued
	Skin reactions	 For signs and symptoms of skin reactions (Including SJS and TEN), exclude other causes 	 Withhold KEYTRUDA for Grade 3 or <u>suspected</u> SJS or TEN, until adverse reactions recover to Grades 0–1* Permanently discontinue KEYTRUDA for Grade 4 or <u>confirmed</u> SJS or TEN For signs or symptoms of SJS or TEN, the patient should be referred to a specialised unit for assessment and treatment
-	Elevated liver enzymes	 For liver enzyme elevations, in patients with RCC being treated with KEYTRUDA in combination with axitinib 	 If ALT or AST ≥3x ULN but <10x ULN without concurrent total bilirubin ≥2x ULN, both KEYTRUDA and axitinib should be withheld until these adverse reactions recover to Grades 0–1. Corticosteroid therapy may be considered If ALT or AST ≥10x ULN or >3x ULN with concurrent total bilirubin ≥2x ULN, both KEYTRUDA and axitinib should be permanently discontinued and corticosteroid therapy may be considered

*If treatment-related toxicity does not resolve to Grades 0–1 within 12 weeks after last dose of KEYTRUDA, or if corticosteroid dosing cannot be reduced to ≤10 mg prednisone or equivalent per day within 12 weeks, KEYTRUDA should be permanently discontinued.





References

- 1. KEYTRUDA Summary of Product Characteristics.
- 2. Harvey, RD. *Clin Pharm Therapeutics* 2014:92(2):214–23.
- 3. Chen DS & Mellman I. *Cell* 2013;39:1–10.
- 4. Bailly C, et al. NAR Cancer 2020; 2(1):doi:10.1093/narcan/zcaa002.
- 5. INLYTA Summary of product characteristics. Available from: <u>https://www.medicines.org.uk/emc/product/4325/smpc.</u> Last updated September 2021.
- 6. Lala M et al. European Journal of Cancer 2020;131:68e75.
- KISPLYX Summary of product characteristics. Available from: <u>https://www.medicines.org.uk/emc/product/7881/smpc</u>. Last updated August 2024.
- 8. PADCEV Summary of product characteristics. Available from: <u>https://www.medicines.org.uk/emc/product/13670/smpc</u>. Last updated November 2024.
- 9. Common Terminology Criteria for Adverse Event (CTCAE). Version 4.0 Published May 2009. Available at https://evs.nci.nih.gov/ftp1/CTCAE/CTCAE_4.03/CTCAE_4.03_2010-06-14_QuickReference_8.5x11.pdf.





Abbreviations

Abbreviation	Definition
5-FU	fluorouracil
AE	adverse event
ALK	anaplastic lymphoma kinase
ALT	alanine aminotransferase
AST	aspartate aminotransferase
BID	twice a day
cHL	classical Hodgkin lymphoma
CPS	combined positive score
EGFR	epidermal growth factor receptor
GVHD	graft-versus-host-disease
HSCT	haematopoietic stem cell transplant
HNSCC	head and neck squamous cell carcinoma
imAE	immune-mediated adverse event
IV	intravenous

Abbreviation	Definition	
NSAIDS	non-steroidal anti-inflammatory drugs	
NSCLC	non-small cell lung carcinoma	
PD-1	programmed death-1	
PD-L1	programmed death ligand-1	
PD-L2	programmed death ligand-2	
QxW	every x weeks	
RCC	renal cell carcinoma	
SIS	Stevens-Johnson syndrome	
TEN	toxic epidermal necrolysis	
TNBC	triple-negative breast cancer	
TPS	tumour proportion score	
ULN	upper limit of normal	
VOD	veno-occlusive disease	





CTCAE Grading Criteria⁹

	Adverse event	Grade 1	Grade 2	Grade 3	Grade 4
3,0-	Adrenal insufficiency	Asymptomatic; clinical or diagnostic observations only; intervention not indicated	Moderate symptoms; medical intervention indicated	Severe symptoms; hospitalisation indicated	Life-threatening consequences; urgent intervention indicated
	ALT increase	>ULN-3.0 x ULN	>3.0–5.0 x ULN	>5.0–20.0 x ULN	>20.0 x ULN
	AST increase	>ULN-3.0 x ULN	>3.0–5.0 x ULN	>5.0–20.0 x ULN	>20.0 x ULN
D -	Colitis	Asymptomatic; clinical or diagnostic observations only; intervention not indicated	Abdominal pain; mucus or blood in stool	Severe abdominal pain; change in bowel habits; medical intervention indicated; peritoneal signs	Life-threatening consequences; urgent intervention indicated
	Hyperthyroidism	Asymptomatic; clinical or diagnostic observations only; intervention not indicated	Symptomatic; thyroid suppression therapy indicated; limiting instrumental ADL	Severe symptoms; limiting self care ADL; hospitalisation indicated	Life-threatening consequences; urgent intervention indicated
	Hyperglycaemia	Fasting glucose value >ULN–160 mg/dL; Fasting glucose value >ULN–8.9 mmol/L	Fasting glucose value >160–250 mg/dL; Fasting glucose value >8.9–13.9 mmol/L	>250–500 mg/dL; >13.9–27.8 mmol/L; hospitalisation indicated	>500 mg/dL; >27.8 mmol/L; life-threatening consequences
	Hypothyroidism	Asymptomatic; clinical or diagnostic observations only; intervention not indicated	Symptomatic; thyroid replacement therapy indicated; limiting instrumental ADL	Severe symptoms; limiting self care ADL; hospitalisation indicated	Life-threatening consequences; urgent intervention indicated

Please note that Common Terminology Criteria for Adverse Events (CTCAE) referred to in the KEYTRUDA Summary of Product Characteristic is version 4.0, version 5.0 is the current version.

UK

ΡΙ

CTCAE Grading Criteria⁹

	Adverse event	Grade 1	Grade 2	Grade 3	Grade 4		
G	Blood and lymphatic system disorders	Asymptomatic or mild symptoms; clinical or diagnostic observations only; intervention not indicated	Moderate; minimal, local or non- invasive intervention indicated; limiting age appropriate instrumental ADL	Severe or medically significant but not immediately life-threatening; hospitalisation or prolongation of existing hospitalisation indicated; limiting self care ADL	Life-threatening consequences; urgent intervention indicated		
() -	Infusion related reaction	Mild transient reaction; infusion interruption not indicated; intervention not indicated	Therapy or infusion interruption indicated but responds promptly to symptomatic treatment (e.g., antihistamines, NSAIDS, narcotics, IV fluids); prophylactic medications indicated for ≤24 hrs	Prolonged (e.g. not rapidly responsive to symptomatic medication and/or brief interruption of infusion); recurrence of symptoms following initial improvement; hospitalisation indicated for clinical sequelae	Life-threatening consequences; urgent intervention indicated		
	Nephritis (acute kidney injury)	Creatinine level increase of >0.3 mg/dL; creatinine 1.5–2.0 x above baseline	Creatinine 2–3 x above baseline	Creatinine >3 x baseline or >4.0 mg/dL; hospitalisation indicated	Life-threatening consequences; dialysis indicated		
() -	Pneumonitis	Asymptomatic; clinical or diagnostic observations only; intervention not indicated	Symptomatic; medical intervention indicated; limiting instrumental ADL	Severe symptoms; limiting self care ADL; oxygen indicated	Life-threatening respiratory compromise; urgent intervention indicated (e.g. tracheotomy or intubation)		
	Skin reactions	Term covers multiple adverse events					



Please note that Common Terminology Criteria for Adverse Events (CTCAE) referred to in the KEYTRUDA Summary of Product Characteristic is version 4.0, version 5.0 is the current version.

UK

ΡΙ

KEYTRUDA[®] (pembrolizumab)

UK prescribing information can be found at: <u>https://www.emcpi.com/pi/33162</u>

Pooled safety data of KEYTRUDA across all indications and AE management can be found in the Summary of Product Characteristics (SmPC)

Refer to KEYTRUDA SmPC and Risk Minimisation Materials (RMM) before prescribing.

Adverse events should be reported. Reporting forms and information can be found at <u>https://yellowcard.mhra.gov.uk/</u> (This links to an external site) or search for MHRA Yellow Card in the Google Play or Apple App Store. Adverse events should also be reported to MSD, UK (Tel: 0208 154 8000).

*By clicking on the links within this slide, you will be taken to an external website. MSD makes no warranties or representations of any kind as to the accuracy, completeness, reliability or usefulness of any information contained in third party sites and shall have no liability for any loss or damage of any kind that may arise from your use of such content or information. Inclusion of any third-party link does not imply an endorsement or recommendation by MSD.

If you have any questions or would like to request any further materials, please contact: MSD medical information (0208 154 8000, <u>medicalinformationuk@msd.com</u>)

Legal Category: POM

Copyright © 2025 Merck & Co., Inc., Rahway, NJ, USA and its affiliates. All rights reserved.