

**KEYTRUDA SC™**  
(pembrolizumab)  
subcutaneous injection

395 mg/2.4 mL  
790 mg/4.8 mL

# Dosing, storage, and preparation of KEYTRUDA SC™ for subcutaneous injection


## Dosing schedules

Two dosing options and solution-strength vials are available for KEYTRUDA SC:<sup>1,2</sup>

**Dose: Q3W**

Administered subcutaneously over **1 minute\***

Low injection volume: 2.4 mL  
**395 mg pembrolizumab**  
**Yellow cap**




OR

**Dose: Q6W**

Administered subcutaneously over **2 minutes\***

Low injection volume: 4.8 mL  
**790 mg pembrolizumab**  
**Green cap**



Vials shown to scale, not actual size.

## Dosing video

[Click here](#) to watch the Dosing, Preparation, and Administration video



## Storage

### Storage of unopened vials:<sup>1</sup>

**KEYTRUDA SC has the same storage requirements as KEYTRUDA IV<sup>1,3</sup>**

- Store in a refrigerator (2°C–8°C)
- Store in the original carton to protect from light
- Do not shake or freeze vials

If removed from refrigeration, unpunctured vials can remain at room temperature (≤25°C) for up to 24 hours before preparation for administration.

### Storage of the prepared syringes:<sup>1</sup>

This product does not contain preservative and **should be used immediately after withdrawing from the vial**. If not used immediately, store the syringe containing KEYTRUDA SC with the transfer needle and cap in place for:<sup>\*\*</sup>

Up to 8 hours at room temperature OR Up to 24 hours in the refrigerator (2°C–8°C)

The 24-hour period may include up to 8 hours at room temperature.

If refrigerated, the filled syringe must be allowed to come to room temperature for at least 30 minutes prior to administration.

**The syringe must not be frozen.**

Do not attach the hypodermic needle until immediately prior to administration to avoid clogging.

### Discard if storage time exceeds these limits

- Do not store any unused portion of the solution for the injection for reuse
- Any unused medicine or waste material should be disposed of in accordance with local requirements

## Preparation


### How to prepare the KEYTRUDA SC syringe for injection:<sup>1</sup>

**1 Check the vial**

Check the vial label to ensure the correct formulation is prepared and administered to the patient as prescribed.

Visually inspect the vial for particulate matter and discoloration. The solution is clear to slightly opalescent, colourless to slightly yellow.

Discard the vial if visible particles or discoloration are observed.




**3 Withdraw the required volume**

Withdraw either 2.4 mL (395 mg) or 4.8 mL (790 mg) using a sterile syringe and a transfer needle (18–21G recommended).

KEYTRUDA SC is compatible with polypropylene and polycarbonate syringe material and stainless-steel transfer and injection needles.

KEYTRUDA SC vial is for single use only. Discard the empty vial or any unused portion left in the vial.\*




Preparation should be performed by trained personnel in accordance with your local hospital guidance and with good practice (e.g. aseptic technique)

**2 Bring the vial to room temperature**

Allow the refrigerated vial to come to room temperature for at least 30 minutes. Prior to the preparation for administration, if needed, the unpunctured vial may be stored at room temperature for up to 24 hours.


Do not dilute, do not shake the vial.



**4 Change to injection needle before use**

To avoid needle clogging, change the needle to a 25–30G, 13 mm hypodermic injection needle immediately prior to subcutaneous injection.

Do not attach the hypodermic needle until immediately prior to administration to avoid clogging.



## Administration


### KEYTRUDA SC is administered over 1 or 2 minutes, with two subcutaneous injection-site options: the abdomen or thigh<sup>1,2</sup>

Should be administered by an HCP and must be administered by subcutaneous injection only:

**Q3W** Inject over **1 minute\*** one dose (395 mg/2.4 mL)

OR

**Q6W** Inject over **2 minutes\*** one dose (790 mg/4.8 mL)



### Administration techniques

- Inject into the subcutaneous tissue of the **thigh or abdomen**, avoiding the 5 cm area around the navel
- Do not inject into skin that is damaged, sore, bruised, scarred, scaly, or has red patches
- Rotate injection sites** for subsequent injections (ensure the injection site is at least 2.5 cm from the previous injection site)
- During treatment with KEYTRUDA SC do not administer other medicinal products for subcutaneous use at the same site
- To best prepare for injection, consider recommending that your patients wear loose-fitting clothes, such as a skirt or shorts

## Disposal

- Vials are for single use only and any unused portion should be discarded
- Dispose in accordance with local requirements



## Prescribing Information

KEYTRUDA SC is licensed for use in adult patients across most KEYTRUDA IV indications, whether alone or in combination with other therapies.<sup>1,3</sup>

One such indication: KEYTRUDA IV and SC, in combination with pemetrexed and platinum chemotherapy, are each indicated for the first-line treatment of metastatic non-squamous non-small cell lung carcinoma in adults whose tumours have no EGFR or ALK positive mutations.<sup>1,3</sup>

Please consult the SmPC and risk minimisation materials for further information before making prescribing decisions.

[Click here](#) for: **KEYTRUDA SC 395 mg/2.4 mL** UK Prescribing Information  
This link will direct you to a third-party website.

[Click here](#) for: **KEYTRUDA SC 790 mg/4.8 mL** UK Prescribing Information  
This link will direct you to a third-party website.

[Click here](#) for: **KEYTRUDA IV** UK Prescribing Information  
This link will direct you to a third-party website.

Adverse events should be reported. Reporting forms and information can be found at <https://yellowcard.mhra.gov.uk/> or search for MHRA Yellow Card in the Google Play or Apple App Store. Adverse events should also be reported to MSD (Tel: 020 8154 8000).

\*Does not account for all aspects of treatment. Actual clinic time may vary; \*\*In-use storage times and conditions prior to use are the responsibility of the user; †Any unused medicinal product or waste material should be disposed of in accordance with local requirements. ALK, anaplastic lymphoma kinase; EGFR, epidermal growth factor receptor; G, gauge; HCP, healthcare professional; IV, intravenous; MHRA, Medicines and Healthcare products Regulatory Agency; Q3W, every 3 weeks; Q6W, every 6 weeks; SC, subcutaneous; SmPC, Summary of Product Characteristics. 1. KEYTRUDA (pembrolizumab) 395 mg Solution for Injection Summary of Product Characteristics; 2. KEYTRUDA (pembrolizumab) 790 mg Solution for Injection Summary of Product Characteristics; 3. KEYTRUDA (pembrolizumab) Solution for Infusion Summary of Product Characteristics.